

PORTLAND HARBOR RI/FS

ROUND 3B FISH AND INVERTEBRATE TISSUE AND COLLOCATED SEDIMENT DATA REPORT

ADDENDUM 1: LEAD AND ANTIMONY RESULTS FOR SMALLMOUTH BASS SAMPLE LW3-SB010E-C00B

DRAFT

DO NOT QUOTE OR CITE

This document is currently under review by US EPA and its federal, state, and tribal partners, and is subject to change in whole or in part.

September 26, 2008

Prepared for

The Lower Willamette Group

Prepared by

Integral Consulting Inc.

IC08-0012

USEPA SF 1290654

Round 3B Tissue and Collocated Sediment Data Report Addendum 1: Lead and Antimony in LW3-SB010E-C00B September 26, 2008 DRAFT

TABLE OF CONTENTS

TABL	E OF CONTENTS	i
	OF TABLES	
	OF ACRONYMS	
	INTRODUCTION	
	LABORATORY ANALYSES	
	DATA VALIDATION AND DATA QUALITY	
	REANALYSIS RESULTS	
	REFERENCES	



Table 4-3.

Round 3B Tissue and Collocated Sediment Data Report Addendum 1: Lead and Antimony in LW3-SB010E-C00B September 26, 2008 DRAFT

LIST OF TABLES

- Table 4-1. Results for Reanalysis of Smallmouth Bass Sample LW3-SB010E-C00B for Lead and Antimony
 Table 4-2. Lead and Antimony Values for Round 3B Bass Samples
 - Revised Lead and Antimony Values for Sample LW3-SB010E-C00B in SCRA Database





Round 3B Tissue and Collocated Sediment Data Report Addendum 1: Lead and Antimony in LW3-SB010E-C00B September 26, 2008 DRAFT

LIST OF ACRONYMS

CAS	Columbia Analytical Services, Inc.
CLP	Contract Laboratory Program
EPA	U.S. Environmental Protection Agency
QC	quality control
QAPP	quality assurance project plan
RI/FS	remedial investigation and feasibility study
SCRA	site characterization and risk assessment



Round 3B Tissue and Collocated Sediment Data Report Addendum 1: Lead and Antimony in LW3-SB010E-C00B September 26, 2008

1.0 INTRODUCTION

Laboratory results for the Round 3B fish tissue samples included anomalously high lead and antimony levels in one smallmouth bass body sample (i.e. the body remaining after removal of the fillets; includes gut and gut contents). The individual bass for this composite sample were collected from the east side of the Willamette River between river miles 9.5 and 10.5. The concentrations of lead and antimony in this sample were greater than 200 times those found in the other smallmouth bass samples from the Portland Harbor study area. However, the results for the corresponding fillet sample were comparable to other Portland Harbor smallmouth bass fillet samples. The body sample, LW3-SB010E-C00B, was reanalyzed to confirm the high lead and antimony concentrations in this sample. Reanalysis procedures and results are presented in this document.

Round 3B Tissue and Collocated Sediment Data Report Addendum 1: Lead and Antimony in LW3-SB010E-C00B September 26, 2008 DRAFT

2.0 LABORATORY ANALYSES

The original analyses for metals in smallmouth bass sample LW3-SB010E-C00B and the reanalyses for lead and antimony were all completed by Columbia Analytical Services (CAS), located in Kelso, Washington. The composite tissue sample was archived frozen at -20°C from the time of collection until reanalysis, except during subsampling for the original analyses, in compliance with quality assurance project plan (QAPP) requirements. The holding time of one year and sample storage requirements were met for the reanalyses.

The sample was thawed, thoroughly mixed, and three subsamples (coded A, B, and C) were collected from three different areas of the sample jar. Each of the subsamples was freeze-dried separately and rehomogenized. The subsamples were digested and analyzed in duplicate (coded A1, A2, B1, etc.). A laboratory duplicate was also prepared from subsample A (coded A1D). Subsample A was therefore analyzed in triplicate and subsamples B and C were analyzed in duplicate, for a total of seven reanalyses.

One laboratory deviation from QAPP Addendum 9 (Integral 2007) was noted. The sample reanalyses were performed by EPA method 6020. The analytical method for lead provided in QAPP Addendum 9 is EPA method 6010B, and this method was used for the original lead analysis. However, EPA method 6020 is commonly used for lead analyses and laboratory quality control (QC) results were acceptable. The use of method 6020 rather than 6010B for lead does not affect the quality or comparability of the lead data in any way.

Round 3B Tissue and Collocated Sediment Data Report Addendum 1: Lead and Antimony in LW3-SB010E-C00B September 26, 2008 DRAFT

3.0 DATA VALIDATION AND DATA QUALITY

The reanalysis data were validated by Integral Consulting Inc. in accordance with guidance provided in U.S. Environmental Protection Agency [EPA] Contract Laboratory Program [CLP] National Functional Guidelines for Inorganic Data Review (EPA 2004) and Guidance on Environmental Data Verification and Validation (EPA 2002), as specified in the Round 2 QAPP.

The results were subjected to full data validation, including the evaluation and assessment of the sample results and applicable quality control results reported by the laboratory and recalculations of laboratory results. Data validation results are summarized as follows:

- Instrument operating conditions were correct and the instrument was properly calibrated.
- Laboratory accuracy was acceptable as demonstrated by the matrix spike recovery for antimony and certified reference material recovery for lead.
- Precision for the various replicates was greater than the control limit of 30%, with 41 relative percent difference reported for the laboratory duplicate for both antimony and lead and relative standard deviations greater than 40% for all replicates for both metals.

All reanalysis data were qualified as estimated based on the variability of the replicate results. The data are usable for all intended purposes, with the knowledge that they may be less precise than unqualified data.

Round 3B Tissue and Collocated Sediment Data Report Addendum 1: Lead and Antimony in LW3-SB010E-C00B September 26, 2008

DRAFT

4.0 REANALYSIS RESULTS

The data for the reanalyses confirm the original results (Table 4-1). For lead, the original result was 2230 mg/kg, and the reanalysis results ranged from 908 to 2590 mg/kg wet weight. For antimony, the original result was 10.9 mg/kg and the reanalysis results ranged from 4.9 to 13.4 mg/kg wet weight.

The anomalous result for this single bass body composite will be evaluated in the draft RI report in the context of lead and antimony concentrations found for other media in this location and throughout the site. A preliminary review of the other data types (e.g., other biota tissues, surface sediments) from the RM 10 east vicinity do not show elevated lead or antimony levels.

Alloys of lead and antimony are widely used in a variety of industrial and recreational equipment. Recreational uses for antimonial lead include fishing gear such as sinkers, lures, and jigs, and shotgun pellets.

The total mass of lead in sample LW3- SB010E-C00B was approximately 2.8 grams¹, or about 0.1 ounce. This quantity is consistent with a small fishing sinker. It seems likely that a fish could have ingested an antimonial lead sinker from a baited fishing lure or jig.

The individual results for the original analysis and all reanalyses were averaged according to rules described in *Portland Harbor Remedial Investigation/Feasibility Study (RI/FS) Technical Memorandum: Guidelines for Data Averaging and Treatment of Non-detected Values for the Round 1 Database* (Kennedy/Jenks et al. 2004) and the averaged values were incorporated into the site characterization and risk assessment (SCRA) database. The final values are 1640 and 8.41 mg/kg for lead and antimony, somewhat lower than the original concentrations, but still much higher than other bass body samples from the Study Area (Table 4-2). Individual and final averaged concentrations are shown in Table 4-1. The final values in the SCRA database, including revised calculated whole body concentrations, are provided in Table 4-3. The updated SCRA database is available on the Portland Harbor Collaboration Portal.

¹ The product of the lead concentration in the fish body sample, 1640 mg/kg wet weight and the combined weight of the individual fish bodies included in the sample, 1725 g, divided by 1,000,000 to convert to grams.

Portland Harbor RI/FS
Round 3B Tissue and Collocated Sediment Data Report
Addendum 1: Lead and Antimony in LW3-SB010E-C00B
September 26, 2008

5.0 REFERENCES

EPA. 2002. Guidance on Environmental Data Verification and Validation. EPA AQ/G-8. U.S. Environmental Protection Agency, Office of Environmental Information, Washington, DC.

EPA. 2004. USEPA Contract Laboratory Program National Functional Guidelines for Inorganic Data Review. EPA 540-R-04-004. Office of Emergency and Remedial Response, U.S. Environmental Protection Agency, Washington, DC.

Integral. 2007. Portland Harbor RI/FS Round 2 Quality Assurance Project Plan Addendum 9: Fish and Invertebrate Tissue and Collocated Sediment for Round 3B. Final. IC07-0020. Prepared for the Lower Willamette Group, Portland, OR. Integral Consulting Inc., Mercer Island, WA.

Kennedy/Jenks, Integral, and Windward. 2004. Portland Harbor RI/FS Technical Memorandum: Guidelines for Data Reporting, Data Averaging, and Treatment of Non-Detected Values for the Round 1 Database. Kennedy/Jenks Consultants, Portland, OR; Integral Consulting Inc., Bellevue, WA; Windward Environmental LLC, Seattle, WA. June 24, 2004.

Table 4-1. Results for Reanalysis of Smallmouth Bass Sample LW3-SB010E-C00B for Lead and Antimony

	Lead	Antimony
Sample ID_	(mg/kg wet weight)	(mg/kg wet weight)
Original analysis		
LW3-SB010E-C00B	2230	10.9
Reanalysis ^a		
LW3-SB010E-C00B-A1	2350	12.7
LW3-SB010E-C00B-A1Db	1550	8.4
LW3-SB010E-C00B-A2	2590	13.4
LW3-SB010E-C00B-B1	958	5.2
LW3-SB010E-C00B-B2	908	4.9
LW3-SB010E-C00B-C1	1150	5.3
LW3-SB010E-C00B-C2	1320	7.1
Average, values included in database ^c	1640	8.41

Notes:

^a Three subsamples of the fish homogenate were individually freeze-dried. Subsamples B and C were analyzed in duplicate and subsample A was analyzed in triplicate (see footnote b).

^b This lab duplicate is an additional aliquot from freeze-dried subsample A and constitues a third analysis of freeze-dried subsample A (i.e., in addition to samples LW3-SB010E-C00B-A1 and -A2).

^c The average was calculated by first averaging each replicate and then averaging these averaged values. These value are included in the SCRA database.

Table 4-2. Lead and Antimony Values for Round 3B Bass Samples

					ta	Remainder data					
Species	Location	CAS Rn	Analyte	Fillet Sample	Detect flag	Result	Units	Remainder Sample	Detect flag	Result	Units
smallmouth bass	SB02E	7439-92-1	Lead	LW3-SB02E-C00F	Y	0.011	mg/kg	LW3-SB02E-C00B	N	0.019 U	mg/kg
mallmouth bass	SB03E	7439-92-1	Lead	LW3-SB02E-C00F	Ϋ́	0.518	mg/kg	LW3-SB03E-C00B	Y	2.28	mg/kg
smallmouth bass	SB03W	7439-92-1	Lead	LW3-SB03W-C00F	Ϋ́	0.045	mg/kg	LW3-SB03W-C00B	N	0.013 U	mg/kg
smallmouth bass	SB04E	7439-92-1	Lead	LW3-SB04E-C01F	Ý	0.056 T	mg/kg	LW3-SB04E-C01B	Y	0.0315 T	mg/kg
smallmouth bass	SB04W	7439-92-1	Lead	LW3-SB04W-C00F	Ϋ́	0.046	mg/kg	LW3-SB04W-C00B	Ý	0.056	mg/kg
mallmouth bass	SB05W	7439-92-1	Lead	LW3-SB05W-C00F	Ý	0.022	mg/kg	LW3-SB05W-C00B	N	0.016 U	mg/kg
mallmouth bass	SB06E	7439-92-1	Lead	LW3-SB06E-C00F	Ϋ́	0.043	mg/kg	LW3-SB05W-C00B	N	0.018 U	mg/kg
mallmouth bass	SB06W	7439-92-1	Lead	LW3-SB06W-C00F	Ý	0.025	mg/kg	LW3-SB06W-C00B	N	0.015 U	mg/kg
mallmouth bass	SB07E	7439-92-1	Lead	LW3-SB07E-C00F	Ϋ́	0.009	mg/kg	LW3-SB07E-C00B	Y	0.034	mg/kg
mallmouth bass	SB07E	7439-92-1	Lead	LW3-SB07W-C00F	Ϋ́	0.01	mg/kg	LW3-SB07W-C00B	Y	0.023	mg/kg
mallmouth bass	SB08E	7439-92-1	Lead	LW3-SB08E-C00F	Ϋ́	0.007	mg/kg	LW3-SB07W-C00B	N	0.023 0.019 U	mg/kg
mallmouth bass	SB08W	7439-92-1	Lead	LW3-SB08W-C00F	Y	0.007	mg/kg	LW3-SB08W-C00B	N	0.019 U	mg/kg
mallmouth bass	SB09E	7439-92-1	Lead	LW3-SB09E-C00F	Y	0.007 0.007 T	mg/kg	LW3-SB09E-C00B	N	0.017 UT	mg/kg
mallmouth bass	SB09W	7439-92-1	Lead	LW3-SB09W-C00F	Y	0.007	mg/kg	LW3-SB09W-C00B	Y	0.045	mg/kg
mallmouth bass	SB010E	7439-92-1	Lead	LW3-SB010E-C00F	Y	0.022	mg/kg	LW3-SB010E-C00B	Ý	1640 JT	mg/kg
mallmouth bass	SB010E SB010W	7439-92-1	Lead	LW3-SB010W-C00F	Ϋ́	0.011	mg/kg	LW3-SB010E-C00B	Ý	9.8	mg/kg
mallmouth bass	SB011E	7439-92-1	Lead	LW3-SB011E-C00F	Ý	0.013	mg/kg	LW3-SB010W-C00B	Ý	0.165	mg/kg
mallmouth bass	SBOILE	7439-92-1	Lead	LW3-SB011W-C00F	Ϋ́	0.023	mg/kg	LW3-SB011W-C00B	Y	0.144	mg/kg
smallmouth bass	SB02E	7440-36-0	Antimony	LW3-SB02E-C00F	N	0.005 UJ	mg/kg	LW3-SB02E-C00B	N	0.006 UJ	mg/kg
mallmouth bass	SB03E	7440-36-0	Antimony	LW3-SB03E-C00F	N	0.005 UJ	mg/kg	LW3-SB03E-C00B	N	0.006 UJ	mg/kg
mallmouth bass	SB03W	7440-36-0	Antimony	LW3-SB03W-C00F	N	0.005 UJ	mg/kg	LW3-SB03W-C00B	N	0.006 UJ	mg/kg
mallmouth bass	SB04E	7440-36-0	Antimony	LW3-SB04E-C01F	N	0.005 UJT	mg/kg	LW3-SB04E-C01B	N	0.005 UJT	mg/kg
mallmouth bass	SB04W	7440-36-0	Antimony	LW3-SB04W-C00F	N	0.005 UJ	mg/kg	LW3-SB04W-C00B	N	0.006 UJ	mg/kg
mallmouth bass	SB05W	7440-36-0	Antimony	LW3-SB05W-C00F	N	0.005 UJ	mg/kg	LW3-SB05W-C00B	N	0.006 UJ	mg/kg
mallmouth bass	SB06E	7440-36-0	Antimony	LW3-SB06E-C00F	N	0.005 UJ	mg/kg	LW3-SB06E-C00B	N	0.006 UJ	mg/kg
mallmouth bass	SB06W	7440-36-0	Antimony	LW3-SB06W-C00F	N	0.005 UJ	mg/kg	LW3-SB06W-C00B	N	0.005 UJ	mg/kg
mallmouth bass	SB07E	7440-36-0	Antimony	LW3-SB07E-C00F	N	0.005 UJ	mg/kg	LW3-SB07E-C00B	N	0.006 U	mg/kg
mallmouth bass	SB07W	7440-36-0	Antimony	LW3-SB07W-C00F	N	0.005 UJ	mg/kg	LW3-SB07W-C00B	N	0.006 U	mg/kg
mallmouth bass	SB08E	7440-36-0	Antimony	LW3-SB08E-C00F	N	0.005 UJ	mg/kg	LW3-SB08E-C00B	N	0.006 U	mg/kg
mallmouth bass	SB08W	7440-36-0	Antimony	LW3-SB08W-C00F	N	0.005 UJ	mg/kg	LW3-SB08W-C00B	N	0.018 U	mg/kg
mallmouth bass	SB09E	7440-36-0	Antimony	LW3-SB09E-C00F	Y	0.005 JT	mg/kg	LW3-SB09E-C00B	Y	0.006 JT	mg/kg
mallmouth bass	SB09W	7440-36-0	Antimony	LW3-SB09W-C00F	N	0.004 UJ	mg/kg	LW3-SB09W-C00B	N	0.005 U	mg/kg
mailmouth bass	SB010E	7440-36-0	Antimony	LW3-SB010E-C00F	N	0.005 UJ	mg/kg	LW3-SB010E-C00B	Y	8.41 JT	mg/kg
mallmouth bass	SB010W	7440-36-0	Antimony	LW3-SB010W-C00F	N	0.005 UJ	mg/kg	LW3-SB010W-C00B	Y	0.052	mg/kg
smallmouth bass	SB011E	7440-36-0	Antimony	LW3-SB011E-C00F	N	0.005 UJ	mg/kg	LW3-SB011E-C00B	N	0.006 U	mg/kg
smallmouth bass	SB011W	7440-36-0	Antimony	LW3-SB011W-C00F	N	0.005 UJ	mg/kg	LW3-SB011W-C00B	N	0.005 U	mg/kg

Notes

- U The material was analyzed for, but was not detected. The associated numerical value is the sample quantitation limit.
- J The associated numerical value is an estimated quantity.
- T The associated numerical value was mathematically derived (e.g., from summing multiple analyte results such as Aroclors, or calculating the average of multiple results for a single analyte). Also indicates all results that are selected for reporting in preference to other available results (e.g., for parameters reported by multiple methods) for the Round 3 data.

DO NOT QUOTE OR CITE

This document is currently under review by US EPA and its federal, state, and tribal partners, and is subject to change in whole or in part.

Lower Willamette Group

Portland Harbor RI/FS

R3B Tissue and Sediment Data Report
Addendum 1
September 26, 2008
DRAFT

Table 4-3. Revised Lead and Antimony Values for Sample LW3-SB010E-C00B in SCRA Database

LocationID	LocationName	X	Y	Z	RiverMile	Task	SampleDate	SampleTin	ne Species	Tissue	Matrix	LabMatr	ix SampleID	ParentSample SampleType
LW3-SB010E-C00	SB010E	7639790.56	694692.56		9.5 - 10.5	B01-01-67B_Biota	9/6/2007	12:06	smallmouth bass	body without fillet	tissue	tissue	LW3-SB010E-C00B	normal
LW3-SB010E-C00	SB010E	7639790.56	694692.56		9.5 - 10.5	B01-01-67B_Biota	9/6/2007	12:06	smallmouth bass	body without fillet	tissue	tissue	LW3-SB010E-C00B	normal
LW3-SB010E-C00	SB010E	7639790.56	694692.56		9.5 - 10.5	B01-01-67B_Biota	9/6/2007	12:06	smallmouth bass	combined fillet and body w/o fillet fractions	tissue	tissue	LW3-SB010E-C00W	normal
LW3-SB010E-C00	SB010E	7639790.56	694692.56		9.5 - 10.5	B01-01-67B_Biota	9/6/2007	12:06	smallmouth bass	combined fillet and body w/o fillet fractions	tissue	tissue	LW3-SB010E-C00W	normal

Lower Willamette Group

Portland Harbor RI/FS
R3B Tissue and Sediment Data Report
Addendum 1

September 26, 2008 DRAFT

Table 4-3. Revised Lead and Antimony Values for Sample LW3-SB010E-C00B in SCRA Database

LocationID	Method	AnalysisDate	AnalysisTime	AnalyteGroup	cas_rn	Analyte	total_or_dissolved	detect_flag	VALUE	Qualifiers	Units	Basis	DVCategory Notes
LW3-SB010E-C00	SW6020	2/5/2008	11:34	Metals	7440-36-0	Antimony	total	Y	8.41	JT	mg/kg	Wet	QA2Cat1
LW3-SB010E-C00	SW6010B	2/5/2008	10:28	Metals	7439-92-1	Lead	total	Y	1640	JT	mg/kg	Wet	QA2Cat1
LW3-SB010E-C00	SW6020	1/22/2008	14:13	Metals	7439-92-1	Lead	total	Y	1100	JT	mg/kg	Wet	QA2Cat1
LW3-SB010E-C00	SW6020	1/22/2008	15:42	Metals	7440-36-0	Antimony	total	Υ	5.9	JT	mg/kg	Wet	QA2Cat1

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY REGION 10

1200 SIXTH AVENUE SEATTLE, WA 98101

TARGET SHEET

7		g document was not imaged. due to the Original being:
•		Oversized
	X	CD Rom
		Computer Disk
		Video Tape
	,	Other:
	•	
**A copy of the docum	ient may he r	requested from the Superfund Records Center.
Treopy of the docum	ione may be i	equested from the superfund records contor.
	Docu	ment Information
Document ID #:		1290654
- File #:		2.7.1-4
Site Name:		PORSF
	ROUND 3E	FISH AND INVERTEBRATE